



V ACTIV MALOSTAB

**SPECIFIC ADSORBENT PRODUCT
FOR PROMOTING MALO-LACTIC FERMENTATION**

COMPOSITION



Purified yeast hulls and highly pure P.V.P.P. (Polyvinylpolypyrrolidone), with high adsorbent power.

CHARACTERISTICS



V ACTIV MALOSTAB is an enological product based on yeast hulls and P.V.P.P., particularly adapted to the specific removal via adsorption of the toxic substances for the bacteria, as exceeding polyphenols and fatty acids with medium and long chains. This is the best solution to prevent any difficulties that may be encountered in the management of the malolactic fermentation.

Cell walls of the yeast (hulls) are constituted of elements characterized of a high adsorbent power towards lipophilic compounds, as fatty acids with 8, 10 and 12 carbon atoms (octanoic, decanoic and dodecanoic acid) and their esters, substances that are secreted also by the yeasts, particularly and in greater quantities if in stress conditions. The P.V.P.P. selected by Enologica Vason for this formula is characterized by a high and specific adsorbent capacity of phenolic substances, acknowledged as active towards bacteria.


APPLICATIONS



V ACTIV MALOSTAB finds its application as malolactic fermentation regulator. With these specific purposes finds full application in some interesting vinification procedures, especially in the case the mean is particularly adverse for lactic bacteria: for example, in presence of high polyphenols' concentrations or in case of a sluggish alcoholic fermentation with high presence of medium and long fatty acids' chains.

In the case of a stopped malolactic fermentation, V ACTIV MALOSTAB has to be used before the bacteria reinoculation and then wait for the precipitation; after racking the mass, an appropriate and specific nutritional integration has to be done, followed then by the add of the bacteria, as specified in the company's procedures.

When using V ACTIV MALOSTAB comply with the relative legal regulations in force.



V ACTIV MALOSTAB

**SPECIFIC ADSORBENT PRODUCT
FOR PROMOTING MALO-LACTIC FERMENTATION**

DIRECTIONS FOR USE



Disperse V ACTIV MALOSTAB in small water or directly into the product to treat, keep stirred for about 30 minutes and then add to the whole mass. Wait for the precipitation and rack.

DOSAGE



As malolactic fermentation regulator and stop prevention: 50g/hl.
In case of reinoculation of lactic bacteria: up to 100 g/hl.

PACKAGING



Poly laminated bags of 500 g.

STORAGE



Store the product in a cool and dry place. Close the bags properly after use.

HAZARD



Based on the current European regulations the product is classified: not hazardous.